



NEWS POLITICS DOGE HEALTH OPINION VIDEOS SHOWS ORIGINALS

HEALTH

# New Study Finds Concerning Evidence of COVID-19 'Vaccine' Shedding

Another COVID 'conspiracy theory' has been proven true.



Published 3 days ago on December 9, 2024  
By **vnninfluencers**



*This article originally appeared on [Courageous Discourse](#) and was republished with permission.*

Guest post by [Nicolas Hulscher, MPH](#)

A new study titled, [Menstrual Abnormalities Strongly Associated with Proximity to COVID-19 Vaccinated Individuals](#), was just published in the *International Journal of Vaccine Theory, Practice, and Research*:

**In Spring 2021, MyCycleStory<sup>SM</sup> launched a secure online survey to which 92.3% of 6049 respondents self-reported menstrual irregularities occurring after the rollout of the COVID-19 injectables. Each respondent served as her own control because prior to the rollout of COVID-19**


### LATEST




FEAT  
26 F  
Revi  
Janu  
You



NEW:  
Dam  
Forc  
Of li




HEAL  
DISI  
Hidi  
Trial




NEW:  
Hidc  
Dist  
Vacc



HEAL  
COV  
Free  
Exte  
202



NEW:  
'USA'  
NYS  
'Per



NEW:  
Marl  
Pop  
Inau



NEW:  
'Mo  
Ovei



vaccination, the vast majority had regular menstrual cycles. A subgroup of 3390 respondents were only *indirectly exposed* to COVID-19 vaccines or the SARS-CoV-2 virus. This subgroup reported 1) being unvaccinated for COVID-19; 2) having had no COVID-19 symptoms; and 3) no positive test for COVID-19, yet a substantial majority of these women, who were only *indirectly exposed* to COVID-19 injectables or COVID-19 infections still had many of the same menstrual abnormalities as the 2659 women who were *directly exposed* to a COVID-19 injection (798), or had COVID-19 symptoms (1347), or tested positive for COVID-19 (514). Generalized linear mixed modeling was used to examine the association (not assuming causation) between abnormal menses experienced after the COVID-19 vaccine rollout by respondents who were only *indirectly exposed* by some degree of proximity to persons. Chi-Square, Student's *t*, Kruskal-Wallis or ANOVA tests were used to assess the statistical significance of the similarities of menstrual irregularities reported by the *directly exposed* and *indirectly exposed* groups.

The mean age of the entire cohort was  $37.8 \pm 0.1$  years. The percentage of the indirectly exposed participants who reported being within 6 feet of a COVID-19 vaccinated person was 85.5%. Of these, 71.7% had irregular menstrual symptoms within one week and 50.1% had irregular menstrual symptoms within  $\leq 3$  days after exposure. When comparing daily proximity to a vaccinated person, the categories of “daily within 6 feet outside the household” versus “seldom/sometimes/daily outside 6 feet” had the highest relative risk at 1.34 ( $p < 0.01$ ) for heavier menstrual bleeding, early menses at more than 7 days early with a relative risk at 1.28 ( $p = 0.03$ ), and extended bleeding for more than 7 days with relative risk at 1.26 ( $p = 0.04$ ). Indirect exposure to COVID-19 vaccinated persons was significantly associated with the likelihood of the onset of menstrual irregularities. This study provides additional data to complement a growing body of evidence raising concerns regarding the safety of mRNA vaccines.

International Journal of Vaccine Theory, Practice, and Research  
**IJVTPR**

**Menstrual Abnormalities Strongly Associated with Proximity to COVID-19 Vaccinated Individuals**

Sue E. Peters PhD<sup>1\*</sup>, Jill Newman MS<sup>2\*</sup>, Heather Ray<sup>3</sup>, James A. Thorp MD<sup>4</sup>, Tiffany Parotto<sup>5</sup>, Brian Hooker PhD<sup>6</sup>, Dan McDyer MD<sup>7</sup>, Leonard Murphy<sup>8</sup>, Raphael B. Stricker MD<sup>9</sup>, Maureen McDonnell BSN<sup>10</sup>, Paul J. Mills PhD<sup>11</sup>, Warren Gieck<sup>12</sup>, Christiane Northrup MD<sup>13</sup>




Figure 2. A bar chart of the frequency percentages of the response categories to the survey question, “How often are you in close proximity with individuals who have been vaccinated? (choose closest option)” The red bars signify the cumulative responses of proximity within 6 feet, whereas the blue bars signify the cumulative responses of proximity outside 6 feet.




Figure 3. A bar chart of the frequency percentages of the response categories to the survey question: “Do you know how many days between being within close proximity to someone who has been vaccinated and when you began showing symptoms?” The purple bars signify the cumulative responses of symptoms onset >1 week, whereas the orange bars signify the cumulative responses of symptoms onset within 1 week of being in close proximity to a vaccinated person.

**Table 4**  
**Relative Risk for Abnormal Menstrual Symptoms by Proximity to Vaccinated Individuals in Women with No Direct Vaccine or SARS-CoV-2 Exposure (n = 3390)**

Abnormal Symptom Experienced for the First Time	Partner/Live with vaccinated person(s)	Daily within 6 feet outside household	Sometimes within 6 feet	Contrasting p-value*
	vs Seldom/Sometimes/Daily outside 6 feet	vs Seldom/Sometimes/Daily outside 6 feet	vs Seldom/Sometimes/Daily outside 6 feet	
	RR [95% CI]	RR [95% CI]	RR [95% CI]	
Heavier menstrual bleeding than usual	1.16 [0.86, 1.55]	<b>1.34 [1.08, 1.65]</b>	<b>1.26 [1.01, 1.57]</b>	<b>0.048</b>
Early menses (>7 days early)	0.72 [0.50, 1.03]	<b>1.28 [1.03, 1.59]</b>	1.09 [0.86, 1.37]	<b>&lt;.001</b>
Extended menstrual bleeding (>7 days)	0.98 [0.70, 1.35]	<b>1.26 [1.01, 1.57]</b>	1.07 [0.85, 1.36]	0.055
Severe cramping and abdominal discomfort	0.99 [0.70, 1.40]	1.23 [0.97, 1.56]	1.05 [0.82, 1.36]	0.127
Heavy menstrual clotting (larger than a dime)	<b>0.64 [0.42, 0.97]</b>	1.13 [0.88, 1.44]	0.92 [0.71, 1.20]	<b>0.011</b>
Spotting between periods	0.76 [0.50, 1.16]	1.22 [0.93, 1.58]	1.12 [0.85, 1.48]	0.062
Decidual cast shedding	1.17 [0.57, 2.42]	1.40 [0.83, 2.34]	1.34 [0.78, 2.29]	0.621
At least 1 of 6 most prevalent abnormal or irregular symptoms	0.90 [0.77, 1.06]	<b>1.16 [1.05, 1.28]</b>	<b>1.13 [1.02, 1.25]</b>	<b>&lt;.001</b>

RR = Relative Risk; CI = Confidence Interval; Significant contrasts at p < .05 are given in bold print.  
 \* p-value from generalized linear mixed models.

**Pierre Kory, MD MPA** ✓  
@PierreKory · Follow

The most puzzling thing we've seen with the vaccine is its ability to "shed" and harm those who never got it. A peer-reviewed study just validated the thousands of shedding reports sent to us. Please share your story so we can untangle this mystery.

pierrekorymedicalmusings.com  
Newly Published Study Shows Shedding Of Covid mRNA Vaccine Products  
A new study found a strong association of new onset menstrual irregularities with "indirect" exposure to Covid vaccines, i.e. being in ...

3:10 PM · Dec 9, 2024

♥ 11.2K    💬 Reply    🔗 Copy link

[Read 792 replies](#)

In layman's terms, the study found that women with daily close proximity (within 6 feet) to vaccinated individuals outside their household had a **34% higher risk** of heavier bleeding, a **28% higher risk** of menstruation starting over 7 days early, and a **26% higher risk** of bleeding lasting more than 7 days.

The **scientific plausibility** for these findings is supported by several key observations, which are discussed in the manuscript:

- 1. Timing Consistency with Shedding Studies:** 68.4% of respondents reported symptoms within one week of being near a vaccinated individual, with 48.6% experiencing symptoms within 3 days or the same day, aligning with the [FDA's guidance](#) on vaccine shedding timelines.
- 2. Prolonged Presence of Vaccine Components:** The detection of mRNA fragments and spike protein in vaccinated individuals' blood for extended periods (up to 187 days) provides evidence of prolonged circulation of potential transmissible components.
- 3. Documented Excretion Pathways:** Established pathways for the excretion of lipid nanoparticles and spike protein, including saliva, sweat, breast milk, and potentially exhalation, provide mechanisms for environmental transmission.
- 4. Alignment with Previous Studies:** Similar menstrual irregularities, such as heavier bleeding and prolonged cycles, have been documented in vaccinated individuals, reinforcing the relevance of these findings in unvaccinated individuals.
- 5. Potential Mechanisms of Action:** The cytotoxic and estrogen-receptor-modulating properties of the spike protein, along with known inflammatory and autoimmune responses to vaccines,

offer biologically plausible mechanisms for these observed effects

The authors concluded, “Our findings suggest possible indirect transmission of ingredients or products of the COVID-19 vaccines, presumably through shedding, from people who received one or more of the COVID-19 injections.”

Why didn't our regulatory agencies conduct shedding studies before mass product rollout? This and many other critical questions demand immediate answers from federal authorities.

**Nicolas Hulscher, MPH**

Epidemiologist and Foundation Administrator, McCullough Foundation

<http://www.mcculloughfnd.org>

Please consider following the [McCullough Foundation](#) and [Nicolas Hulscher](#) on X (formerly Twitter) for further content.

Copyright 2024 [Courageous Discourse](#)

## Stay Vigilant with VNN.

Get the uncensored truth sent straight to your inbox daily.

RELATED TOPICS: [#COVID-19 VACCINE SHEDDING](#) [#COVID-19 VACCINES](#) [#FEATURED](#)


**DONT MISS**  
CHILLING: Biden EPA Makes First-Ever 'Climate Change' Arrest

**UP NEXT**  
Judge Jeanine Pirro Sets the Record Straight About Daniel Penny and Jordan Neely

### YOU MAY LIKE




**DISTURBING: Pfizer Accused of Hiding Deaths in COVID-19 Vaccine Trials**



**Hidden Pfizer Report Reveals Disturbing Reality for the Vaccinated**



**COVID Vaccine Makers Get Another Free Pass as Biden Administration Extends Liability Shield Through 2029**



**Australian Medical Icon Breaks Down**



**Alarming Levels of DNA**



**New Study Calls for an Immediate**

*TRENDING NOW*



NEWS / 3 days ago

**BUSTED: Taylor Lorenz Says the Quiet Part Out Loud**




NEWS / 2 days ago

**Shocking New Details Emerge About Trump's Would-Be Assassin**



FEATURED / 17 hours ago

**Rep. Thomas Massie Drops Devastating Accusations Against the US Government Over Syria**



NEWS / 2 days ago

**180 Dead: Ha**

